



Merriel, S. W. D., Nadarzynski, T., Kesten, J., Flannagan, C., & Prue, G. (2018). 'Jabs for the boys': time to deliver on HPV vaccination recommendations. *British Journal of General Practice*, 68(674), 406-407. <https://doi.org/10.3399/bjgp18X698429>

Peer reviewed version

Link to published version (if available):
[10.3399/bjgp18X698429](https://doi.org/10.3399/bjgp18X698429)

[Link to publication record in Explore Bristol Research](#)
PDF-document

This is the author accepted manuscript (AAM). The final published version (version of record) is available online via RCGP at <https://bjgp.org/content/68/674/406> . Please refer to any applicable terms of use of the publisher.

University of Bristol - Explore Bristol Research

General rights

This document is made available in accordance with publisher policies. Please cite only the published version using the reference above. Full terms of use are available:
<http://www.bristol.ac.uk/red/research-policy/pure/user-guides/ebr-terms/>

‘Jabs for the boys’ in England – Time to deliver on HPV vaccination recommendations

Dr Samuel W D Merriel¹, Dr Tom Nadarzynski², Dr Joanna M Kesten^{3,4,5}, Dr Carrie Flannagan⁶, Dr Gillian Prue⁷

¹ Centre for Academic Primary Care, Bristol Medical School, University of Bristol, Bristol BS8 2PS, UK

² Department of Psychology, University of Southampton, Southampton SO17 1BJ, UK

³ Bristol Medical School, University of Bristol, Bristol BS8 2PS, UK

⁴ The National Institute for Health Research Health Protection Research Unit in Evaluation of Interventions, Bristol Medical School, University of Bristol, Bristol BS8 2PS, UK

⁵ The National Institute for Health Research Collaboration for Leadership in Applied Health Research and Care West (NIHR CLAHRC West), University Hospitals Bristol NHS Foundation Trust, Bristol BS1 2NT, UK

⁶ Institute of Nursing and Health Research, Ulster University, Londonderry BT52 1SA, UK

⁷ School of Nursing and Midwifery, Queens University, Belfast BT7 1NN, UK

Human papillomavirus (HPV) vaccination programmes in the UK were established in 2008, and targeted females aged 12-13 years. This decision was based on recommendations from the Joint Committee for Vaccination and Immunisation (JCVI), after assessing the available evidence for the impact and cost effectiveness of HPV vaccination programmes. The theory was that reducing HPV rates in females would result in lower infection rates in males of a similar age. This herd immunity did not extend to men-who-have-sex-with-men (MSM) or heterosexual men having sex abroad. Public Health England undertook a trial of a targeted HPV vaccination programme for MSM in 2016/17. The 12-month pilot involved MSM up to 45 years of age being offered *Gardasil 4*, a quadrivalent HPV vaccine, through genitourinary medicine (GUM) and HIV clinics across England. GPs were not included in the pilot as there was insufficient evidence of acceptability and effective delivery. The pilot was reported as a success, and PHE has recommended rolling out this programme across England(1). The JCVI issued an interim statement regarding extending HPV immunisation to adolescent males in July 2017, again concluding that there was insufficient

evidence for benefit in males(2). However, after strong responses through stakeholder consultations, continued pressure from lobby groups such as HPV Action (www.hpvaction.org), and further analysis of the latest research, the JCVI recently recommended extending the HPV vaccination programme to adolescent males on July 18 2018(3).

Human papillomavirus

HPV is a major cause of infection-related cancers. It is associated with cervical, anal, oropharyngeal, penile, vulval and vaginal cancers, as well as anogenital warts. HPV subtypes 16 and 18 are the most prevalent forms, and they are also the subtypes most strongly associated with HPV-related cancers(4). Risk factors for HPV infection include age, number of sexual partners, co-infection with other sexually transmitted diseases, condom use, anal sex, immunosuppression, and cigarette smoking(5). HPV infection rates in women and heterosexual men of corresponding age in England and Wales are expected to fall following the introduction of a HPV vaccination programme for girls in 2008.

The incidence of HPV-related cancers that affect males, such as anal, penile and oropharyngeal cancers, are on the rise. Rates of anal squamous cell carcinoma in England between 1990-2010 in men increased from 0.43 per 100,000 population to 0.73 per 100,000, a rise of 69%(6). HIV-negative MSM have a four-fold higher risk of anal cancer when compared to HIV-negative heterosexual men and women; HIV-positive men have an up to 80-fold higher risk(7). In the US, men were found to have rates of oropharyngeal cancer of 7.6 per 100,000, four times higher than women (1.7 per 100,000) between 2008 -2012. Anogenital warts are caused by a number of HPV strains, not just those covered by currently available HPV vaccines. The total number of patients with genital warts in England from 1971-2001 increased eight-fold, presumed to be mainly due to changing sexual practices(4), although there has been a reduction in incidence in recent years.

HPV vaccination programmes

All nations of the UK currently employ HPV vaccine programmes for adolescent females age 12-13, using the quadrivalent HPV vaccine. Scotland also offers HPV vaccination for MSM through sexual health and HIV clinics.

Forty-two sexual health and HIV clinics in England were included in the pilot programme of vaccinating MSM, with 18,875 MSM attending in that time. 8,580 (45.5%) of MSM attending the clinics in the pilot accepted at least one dose of the vaccine. The vast majority of MSM who completed a feedback survey during their clinic visit (7821/9584 [94.6%]) stated they would prefer to receive subsequent vaccine doses at the same clinic. 677 (7.2%) would have preferred to receive subsequent doses at their GP practice. Only 636 (3.4%) declined the vaccine, meaning the vaccination status of over half of the participants (9659 [51.2%]) in the pilot is unknown.

Internationally, various approaches have been adopted to protect those at high-risk of HPV-related diseases. Australia and the Canadian province of Quebec offer gender-neutral HPV vaccination programmes. Targeted HPV vaccination for MSM under the age of 26 years is available in the US and Ireland. Denmark recently announced a time-limited HPV vaccination of MSM via their GPs. There is on-going debate about the cost effectiveness of gender-neutral versus female-only HPV vaccination, and the efficacy of relying on herd immunity to protect males from HPV infection. Ethical and equality concerns about denying HPV vaccination to an individual on the basis of their gender have also been raised(3).

Delivering an effective HPV vaccination programme

Following the JCVI's recommendation to extend HPV vaccination to males, the Welsh and Scottish governments confirmed they would be implementing the JCVI's advice in offering HPV vaccination to adolescent males as well as females. At the time of writing this editorial, no official response from the English government or the Northern Ireland assembly had been made, and there has been no discussion of any form of catch-up vaccination programme for older males.

HPV vaccination uptake amongst adolescent females in the UK is currently above 80%, with 83.1% of 12-13 year-old females completing a two-dose schedule in 2016/17. This rate has slightly dropped from 86.7% in 2013/14(8). Coverage in the UK remains high despite a significant drop in recent years in coverage in some European countries, such as Denmark and the Republic of Ireland. UK nations that follow the JCVI recommendation and offer HPV vaccination to adolescent

males will need to clearly inform parents of the effects of HPV vaccination for their sons to try to ensure uptake is high enough for the vaccination programme to be effective.

The question needs to be asked whether rolling out a targeted HPV vaccination programme for MSM through sexual health clinics in England alongside a gender-neutral programme for adolescent males and females is an effective use of public health resources. The pilot failed to achieve 50% coverage for the first dose of the HPV vaccine for MSM, and even lower coverage for subsequent doses. The evidence is clear that HPV vaccination is most efficacious when delivered prior to exposure to the virus(9). It is also known that MSM are often reluctant to declare their sexual orientation to healthcare professionals and MSM seldom attend healthcare services prior to sexual debut(10), creating doubts about the effectiveness of a targeted programme delivered via sexual health clinics. Expanding access for a targeted programme to include GPs could boost coverage amongst MSM, however GPs are less confident in identifying young MSM who may benefit compared to sexual healthcare professionals(11).

The International Papillomavirus Society recently released a statement calling for ‘the policy, the resources, political will and the public’s determination’ to move towards eliminating HPV as a public health problem(12). The JCVI recommendation to extend HPV vaccination to males in the UK is the first step in changing policy. Public support for HPV vaccination for males has been strong. Now the political will and proper resources are needed to make gender-neutral HPV vaccination a reality and reduce the incidence of HPV-related diseases for future generations in the UK.

Declarations

SWDM is funded by an NIHR Academic Clinical Fellowship. JMK is partly funded by National Institute for Health Research (NIHR) Collaboration for Leadership in Applied Health Research and Care West (CLAHRC West) at University Hospitals Bristol NHS Foundation Trust and NIHR Health Protection Research Unit in Evaluation of Interventions. The views expressed are those of the authors and not necessarily those of the NHS, the NIHR or the Department of Health and Social Care.

References

1. PHE. Human papillomavirus (HPV) vaccination for Men who have sex with Men (MSM). London; 2018.
2. JCVI. JCVI Interim Statement on Extending HPV Vaccination to Adolescent Boys [Internet]. Public Health England; 2017. p. 1–21. Available from: https://www.gov.uk/government/uploads/system/uploads/attachment_data/file/630125/Extending_HPV_Vaccination.pdf
3. JCVI. Statement on HPV vaccination. London: Joint Committee on Vaccination and Immunisation; 2018.
4. Forman D, De Martel C, Lacey CJ, Soerjomataram I, Lortet-Tieulent J, Bruni L, et al. Global Burden of Human Papillomavirus and Related Diseases. Vaccine [Internet]. 2012 Nov 20;30:12–23. Available from: <http://dx.doi.org/10.1016/j.vaccine.2012.07.055>
5. King E, Gilson R, Beddows S. Human papillomavirus DNA in men who have sex with men: type-specific prevalence, risk factors and implications for vaccination strategies. Br J ... [Internet]. 2015 Mar 19;112(9):1585–93. Available from: <http://www.nature.com/bjc/journal/v112/n9/abs/bjc201590a.html>
6. Wilkinson JR, Morris EJA, Downing A, Finan PJ, Aravani A, Thomas JD, et al. The rising incidence of anal cancer in England 1990-2010: A population-based study. Color Dis [Internet]. 2014 Jun 12;16(7):O234–9. Available from: <http://doi.wiley.com/10.1111/codi.12553>
7. van der Zee R, Richel O, de Vries H, Prins J. The increasing incidence of anal cancer: can it be explained by trends in risk groups? Neth J Med [Internet]. 2013 Oct 14;71(7):401–11. Available from: <http://www.njmonline.nl/getpdf.php?id=1355>
8. PHE. Human Papillomavirus (HPV) vaccination coverage in adolescent females in England: 2016/17 [Internet]. Public Health England; 2017. Available from: www.gov.uk/phe%0Awww.facebook.com/PublicHealthEngland
9. Cutts FT, Franceschi S, Goldie S, Castellsagué X, De Sanjosé S, Garnett G, et al. Human papillomavirus and HPV vaccines: a review. Bull World Health Organ [Internet]. 2007 [cited 2018 Feb 22];85(9):649–732. Available from:

<http://www.who.int/bulletin/volumes/85/9/06-038414/en/#.Wo6OGUkxIDo.mendeley>

10. Nadarzynski T, Smith H, Richardson D, Bremner S, Llewellyn C. Men who have sex with men who do not access sexual health clinics nor disclose sexual orientation are unlikely to receive the HPV vaccine in the UK. *Vaccine* [Internet]. 2018;36(33):5065–70. Available from: <https://doi.org/10.1016/j.vaccine.2018.06.075>
11. Merriel SWD, Flannagan C, Kesten JM, Shapiro GK, Nadarzynski T, Prue G. Knowledge and attitudes of general practitioners and sexual health care professionals regarding human papillomavirus vaccination for young men who have sex with men. *Int J Environ Res Public Health*. 2018;15(1):151.
12. IPVS. IPVS Statement Moving towards Elimination of Cervical Cancer as a Public Health Problem. International Papillomavirus Society; 2018.